

NOTE

A NOTE ON SCHNORR'S SEPARATEDNESS

Daniel J. LEHMANN

Department of Computer Science, University of Warwick, Coventry CV4 7AL, England

Abstract. It is shown that the two concepts of separatedness proposed by Schnorr are related: separatedness implies 1-separatedness. Schnorr's results on 1-separatedness then imply his results on separatedness.

In [1] Schnorr proposed the following two definitions (the notations are those of [1]).

Definition 3.1. Let $f \in \Omega$, then $B \subset \text{mon}(f)$ is called *separated* iff $\forall r \in \text{mon}(f) : \forall s, t \in B : r \geq s \cdot t \Rightarrow [r = s \text{ or } r = t]$.

Definition 5.1. A subset $B \subset \text{mon}(f)$ is called *1-separated* if (S1), (S2) hold.

(S1) $\forall r \in \text{mon}(f) : \forall s, t \in B : r \geq s \cdot t \Rightarrow [r \leq s \text{ or } r \leq t]$,

(S2) $\forall r \in \text{mon}(f) : \forall s \in B : r \geq s \Rightarrow r = s$.

We claim:

Theorem 1. (S1) \Rightarrow (S2).

Proof. Suppose $r \in \text{mon}(f)$, $s \in B$ and $r \geq s$ then $r \geq s \cdot s$ and by (S1) $r \leq s$. But $r \geq s$ and $r \leq s$ imply $r = s$. As a consequence, the condition (S2) can be dropped in the definition of 1-separatedness. \square

Theorem 2. If a subset $B \subset \text{mon}(f)$ is separated then it is 1-separated.

Proof. Obvious. \square

Theorem 3. Let $f \in \Omega$, $\bar{\#}_1(f) \geq \bar{\#}(f)$ and $\#_1(f) \geq \#(f)$.

It follows that Schnorr's Theorem 5.2 implies his Theorem 3.2. Lemmas 3.4, 3.5 and 3.6 can be dispensed with.

Reference

- [1] C.P. Schnorr, A lower bound on the number of additions in monotone computations, *Theoretical Computer Sci.* 2 (1976) 305-315.